

Cephalometric Norms for Saudi Adults Living in the Western Region of Saudi Arabia

Ali H. Hassan^a

Abstract: European-American norms are still used in the orthodontic treatment of Saudi patients, despite the different ethnic backgrounds of Saudis. The aims of this study were to evaluate the cephalometric features of a Saudi population and to establish cephalometric norms for Saudis living in the western region of Saudi Arabia. Seventy lateral cephalometric radiographs of Saudis (32 females and 38 males; aged 18–28 years) with acceptable profiles and Class I dental relationships were traced and analyzed. The mean value, standard deviation, and range of 16 angular and linear variables were calculated. The resulting norms for Saudis were compared with European-American norms using an independent *t*-test. Male and female groups were also compared using the *t*-test. Saudis tend to have an increased ANB angle because of retrognathic mandibles and bimaxillary protrusion as compared with European-Americans. Males tend to have more prognathic mandibles than females as indicated by the statistically significant increase in facial angle ($P < .05$) and SNB angle ($P < .05$). Although the anterior lower face height was similar in males and females, males tend to have a steeper mandibular plane angle when related to the anterior cranial base than females ($P < .05$). Saudis have distinct cephalometric features, which should be used as a reference in treating Saudi orthodontic patients. (*Angle Orthod* 2006;76:109–113.)

Key Words: Saudi Norms; Cephalometric; Norms

INTRODUCTION

Orthodontic treatment is best when the facial and cephalometric characteristics of the ethnic background of patients are considered. The orthodontic literature contains many studies involving cephalometric and profile standards of European-American, African-American, Japanese, and Chinese populations^{1–17} but little for Arabs and Saudis in specific.^{18–24}

Bishara et al¹⁹ established cephalometric standards for Egyptian adolescent boys and girls and compared them with a matched Iowa adolescent sample. There was a great similarity in the overall facial morphology between the Egyptian and Iowan populations. Hamdan and Rock²¹ evaluated the cephalometric features of a Jordanian population as compared with the Eastman standards and found different skeletal and dental

cephalometric features for the Jordanians. Shalhoub et al²² evaluated lateral cephalometric radiographs of 48 adult Saudis with normal facial proportions, compared them with a North American sample, and established a set of cephalometric norms for Saudi adults living in Riyadh. Sarhan and Nashashibi²³ compared cephalometric radiographs of Saudi boys (10–14 years old) with a similar British sample. They found slightly more prognathic Saudi faces, more protruded incisors and lower gonial and saddle angles as compared with the British sample. Al-Jasser²⁴ described the craniofacial characteristics of 87 Saudi students with acceptable profiles and occlusions and compared them with Steiner's European-American standards. It was also concluded that Saudis have different craniofacial features when compared with Steiner norms.

Unfortunately, all the previously mentioned studies were performed in the central region of Saudi Arabia and do not represent the multiracial background of the Saudis. In addition, no single study has been performed in the other regions of Saudi Arabia. The western region of Saudi Arabia, also known as Hijaz, is unique in its ethnic diversity that is mainly because of the Hajj, where Muslims from all over the world come to attend this yearly Islamic pilgrimage in Makkah. Saudis who live in this region are of mixed ethnic origin

^a Assistant Professor, Department of Preventive Dental Sciences, Faculty of Dentistry, King Abdulaziz University, Jeddah, Saudi Arabia.

Corresponding author: Ali H. Hassan, BDS, PhD, Department of Preventive Dental Sciences, Faculty of Dentistry, King Abdulaziz University, PO Box 122423, Jeddah 21332, Saudi Arabia (e-mail: alireserach@gawab.com).

Accepted: January 2005. Submitted: October 2004.
© 2006 by The EH Angle Education and Research Foundation, Inc.




FIGURE 1. The different cephalometric landmarks used.

TABLE 1. Different Linear and Angular Measurements Used

N-PG-FH	Intersection between N-PG and Frankfort horizontal (FH) plane
SN-FH	Angle between SN plane and FH plane
SNA	Maxillary apical base relationship to anterior cranial base
SNB	Mandibular apical base relationship to anterior cranial base
ANB	Apical base relationship
NA-APg	Angle of convexity
FMA	Inclination of mandibular plane to FH
MP-SN	Inclination of mandibular plane angle to anterior cranial base
OC-PL-SN	Inclination of occlusal plane to anterior cranial base
y-axis	Angle made between SN and NGn line
L-FC-HT	Lower face height (anterior nasal spine-Menton)
N-S-BA	Cranial base angle
U1-SN	Inclination of maxillary incisors to anterior cranial base
U1-NA (angle)	Inclination of maxillary incisors to NA
U1-NA (mm)	Protrusion of maxillary incisors to NA
U1-L1	Inclination of maxillary incisors to mandibular incisors
L1-MP	Inclination of mandibular incisors to mandibular plane
L1-NB	Inclination of mandibular incisors to NB
L1-NB (mm)	Protrusion of maxillary incisors to NB

and descendants of Arabs, Indians, Turks, Indonesians, Africans and others. Most of them settled in the western region and eventually became Saudis.

Vorhies and Adams²⁵ simplified the reading of Downs' cephalometric norms, when they developed a polygon or wiggle, in which cephalometric readings were expressed graphically. A wiggle, as described by Vorhies and Adams,²⁵ is a graph in which all average norms are plotted on a central vertical line. The maximum and the minimum readings of each norm are plotted on either side of the central line in a manner that all the Class II readings are placed on the left side

and the Class III readings are placed on the right side of the central line.²⁵

The objectives of this study were to evaluate the cephalometric features of a Saudi population living in the western area of Saudi Arabia, to establish Saudi norms in this area, and to present them diagrammatically in the form of a polygon for easier use.

MATERIALS AND METHODS

A total of 70 lateral cephalometric radiographs of Saudi adults (32 females and 38 males; aged 18–28




FIGURE 2. Wiggle for Saudi Norms.

years) with balanced and acceptable facial profiles, minimum overbite and overjet, Class I skeletal and dental relationships, and no previous orthodontic treatment were traced and analyzed manually by a single examiner. All selected subjects were Arab Saudis (by nationality) living in the western region of Saudi Arabia. A total of 16 angular and linear measurements were calculated (Figure 1; Table 1). The mean value, standard deviation, and range for each variable were calculated. Measurements were compared with European-American norms, and the differences were analyzed and highlighted. European-American norms were derived from Downs,^{1,2} Hasund,²⁶ and Riedel³ analyses. Independent sample *t*-tests were performed to compare Saudis and European-Americans and to compare male and female groups.²⁷ To assess tracing errors, a second tracing was prepared for each of 10 tracings. The mean error in linear measurements was 0.45 mm and in angular measurements was 0.93°.

A set of cephalometric values for male and female Saudis was established. The resulting data (means and standard deviations) were represented diagrammatically in the form of a polygon (wiggle), using the program Microsoft Excel (Microsoft, Redmond, Wash) (Figure 2). The standard deviations were used instead of the maximum and minimum readings in the polygon, unlike the classic wiggle of Vorhies and Adams.²⁵

RESULTS

Compared with European-Americans, Saudis were found to have an increased facial convexity (ANB, 3.65 ± 1.65), a more convex profile (NA-APg = 4 ± 3.5), and a steeper mandibular plane (FMA, 28.0 ± 5.8). In addition, the upper and lower incisors were significantly more proclined and more protruded. The y-axis angle was significantly steeper ($P < .001$) and the anterior lower face height insignificantly shorter in Saudis than in European-Americans (Table 2).

Males were found to have more prognathic mandibles than female as indicated by the statistically significant increase in SNB ($P < .05$). Although the anterior lower face height was similar in males and females, males tended to have a steeper mandibular plane angle than females when related to the anterior cranial base ($P < .05$) (Table 3).

DISCUSSION

The racial, facial, and skeletal characteristics of the patient play a critical role in orthodontic treatment planning. The objectives of this study were to evaluate the cephalometric features of a Saudi population and to establish norms for the Saudis living in the western region of Saudi Arabia. This study was performed using a relatively larger sample size from the Saudi population than those used in previous studies. In addi-

TABLE 2. Saudi-adult Standards as Compared with European-American Standards Using *t*-test

Variables	Saudis (Adults) (n = 70)		European-American(s) (Adults) (n = 48-93)		<i>t</i>	<i>P</i>
	Mean	SD	Mean	SD		
N-PG-FH	86.6	3.64	87.8	3.57	0.98	>.05
SNA	80.8	4.06	82.01	3.89	0.85	>.05
SNB	77.5	4.48	79.97	3.69	1.69	>.05
ANB	3.7	1.522	2.04	1.81	2.59	<.05
NA-APg	5.01	3.05	1.62	4.78	2.8	<.01
Mandibular plane/FH	28.5	4.79	22.4	5.6	3.57	<.001
MP-SN	35.9	5.96	31.7	5.19	2.12	<.05
y-axis	69.6	4.2	59.4	3.82	7.16	<.001
U1-SN	107.8	8.07	103.97	5.75	1.46	>.05
U1-NAZ	27.3	7.5	22	6	2.13	<.05
U1-NA (mm)	6.8	2.9	6	1.9	1.02	>.05
U1-L1	120.6	11.89	130.9	9.24	2.64	<.01
L1-MP	93.9	7.7	93.09	6.78	0.32	>.05
L1-NB	29.34	6.89	25	6	1.87	>.05
L1-NB (mm)	7.52	2.63	5	1.7	2.08	<.05
L-FC-HT (%)	56.03	2.7	57	Not specified		

TABLE 3. Comparison of the Measurements Between Saudi Males and Females Using *t*-test

Variable	Male		Female		<i>P</i>
	Mean	SD	Mean	SD	
N-PG-HF	87.2	4.4	83.5	3.7	<.05*
SNA	80.9	4.9	79.6	3.9	>.05
SNB	77.2	5	75.5	3.5	<.05*
ANB	3.7	3	4.1	2.9	>.05
NA-APg	6.9	6.5	8.9	11.1	>.05
FMA	25.8	5.7	27.8	5.3	>.05
MP-SN	35.4	6.2	33.2	5.5	<.05*
OC-PL-SN	13.3	9.3	20.5	6.5	>.05
y-axis	70.2	6.2	70.3	6.4	>.05
U1-SN	107.9	8.8	103.9	8.8	>.05
U1-NA (angle)	23.5	8.1	23.1	8.4	>.05
U1-NA (mm)	7.1	3.4	6.6	3.4	>.05
U1-L1	120.3	11.3	117.4	12.5	>.05
L1-MP	97.5	9.6	96.6	9.3	>.05
L1-NBZ	29.5	7.3	30.7	7.3	>.05
L1-NB (mm)	7.2	2.9	7.5	3	>.05
L-FC-HT	55.8	3.3	54.8	2.8	>.05

tion, the sample was selected carefully to include Saudis, by nationality, who had Class I skeletal and dental relationships and pleasant faces. By definition, the Saudi population, especially in the western area, is a multiracial mixed population that consists of people who have lived in Saudi Arabia for a long period of time and eventually obtained nationality. Because we believe that Saudis living in this area represent the new Saudi race, which has been established because of interbreeding among the different communities, selection was open to include all Saudi Arabs living in the western province of Saudi Arabia.

Results are consistent with previous studies in Riyadh, Central Province, in that Saudis tend to have

bimaxillary protrusion. Important findings are the increased ANB angle and mandibular plane angle in Saudis as compared with European-Americans with Class I skeletal relationship, which adds more value for using these measurements in evaluating skeletal relationships in Saudis (Table 2).

The polygon is considered a versatile tool for practical clinical use, which simplifies the reading and its presentation to the patients. In addition, presenting the norms of such a mixed race of Saudis is advantageous in the sense that it counts for the expected variability by including norms within one standard deviation.

CONCLUSION

Saudis have distinct cephalometric features, for which specific norms should be used as a reference in treating orthodontic patients. In addition, presenting Saudi norms on a polygon is a faster and practical method of analyzing cephalometrics.

ACKNOWLEDGMENT

The authors acknowledge Dr Budi Kusnoto, Clinical Assistant Professor of Orthodontics, University of Illinois, for his kind help in developing the wiggle.

REFERENCES

- Downs WB. Analysis of the dentofacial profile. *Angle Orthod.* 1956;26:191.
- Downs WB. Variation in facial relationships: their significance in the treatment and prognosis. *Am J Orthod.* 1948; 34:812-840.
- Riedel RA. The relation of maxillary structures to cranium in malocclusion and in normal occlusion. *Angle Orthod.* 1952;22:142-145.
- Steiner CC. The use of cephalometrics as an aid to planning

- and assessing orthodontic treatment. *Am J Orthod.* 1960; 29:8.
5. Bishara SE. Longitudinal cephalometric standards from 5 years of age to adulthood. *Am J Orthod.* 1981;79:35–44.
 6. Scheideman GB, Bell WH, Legan HL, Finn RA, Reisch JS. Cephalometric analysis of dentofacial-normals. *Am J Orthod.* 1980;78:404–420.
 7. Argyropoulos E, Sassouni V. Comparison of the dentofacial patterns for native Greek and American-Caucasian adolescents. *Am J Orthod Dentofacial Orthop.* 1989;95:238–249.
 8. Franchi L, Baccetti T, McNamara JA Jr. Cephalometric floating norms for North American adults. *Angle Orthod.* 1998; 68:497–502.
 9. Farkas LG, Tompson B, Phillips JH, Katic MJ, Cornfoot ML. Comparison of anthropometric and cephalometric measurements of the adult face. *J Craniofac Surg.* 1999;10:18–25.
 10. Huang WJ, Taylor RW, Dasanayake AP. Determining cephalometric norms for Caucasians and African Americans in Birmingham. *Angle Orthod.* 1998;68:503–512.
 11. Anderson AA, Anderson AC, Hornbuckle AC, Hornbuckle K. Biological derivation of a range of cephalometric norms for children of African American descent (after Steiner). *Am J Orthod Dentofacial Orthop.* 2000;118:90–100.
 12. Bailey KL, Taylor RW. Mesh diagram cephalometric norms for Americans of African descent. *Am J Orthod Dentofacial Orthop.* 1998;114:218–223.
 13. Miyajima K, McNamara JA Jr, Kimura T, Murata S, Iizuka T. Craniofacial structure of Japanese and European-American adults with normal occlusions and well-balanced faces. *Am J Orthod Dentofacial Orthop.* 1996;110:431–438.
 14. Engel G, Spolter BM. Cephalometric and visual norms for a Japanese population. *Am J Orthod.* 1981;80:48–60.
 15. Kondo S, Wakatsuki E, Shibagaki H. A somatometric study of the head and face in Japanese adolescents. *Okajimas Folia Anat Jpn.* 1999;76:179–185.
 16. Alcalde RE, Jinno T, Pogrel MA, Matsumura T. Cephalometric norms in Japanese adults. *J Oral Maxillofac Surg.* 1998;56:129–134.
 17. So LL, Davis PJ, King NM. “Wits” appraisal in Southern Chinese children. *Angle Orthod.* 1990;60:43–48.
 18. Loutfy MS, Ponitz P, Harris JE. Cephalometric standards for normal Egyptian face. *Egypt Dent J.* 1971;17:91–100.
 19. Bishara SE, Abdalla EM, Hoppens BJ. Cephalometric comparisons of dentofacial parameters between Egyptian and North American adolescents. *Am J Orthod Dentofacial Orthop.* 1990;97:413–421.
 20. Mouakeh M. Cephalometric evaluation of craniofacial pattern of Syrian children with Class III malocclusion. *Am J Orthod Dentofacial Orthop.* 2001;119:640–649.
 21. Hamdan AM, Rock WP. Cephalometric norms in an Arabic population. *J Orthod.* 2001;28:297–300.
 22. Shalhoub SY, Sarhan OA, Shaikh HS. Adult cephalometric norms for Saudi Arabians with a comparison of values for Saudi and North American Caucasians. *Br J Orthod.* 1987; 14:273–279.
 23. Sarhan OA, Nashashibi IA. A comparative study between two randomly selected samples from which to derive standards for craniofacial measurements. *J Oral Rehabil.* 1988; 15:251–255.
 24. Al-Jasser NM. Cephalometric evaluation of craniofacial variations in normal Saudi population according to Steiner analysis. *Saudi Med J.* 2000;21:746–750.
 25. Vorhies JM, Adams JW. Polygonic interpretation of cephalometric findings. *Angle Orthod.* 1951;21:194–197.
 26. Bosch C, Athanasiou AE. Landmarks, variables and norms of various numerical cephalometric analyses—Cephalometric morphologic and growth data references. In: Athanasiou AE. *Orthodontic Cephalometry.* London, England: Mosby-Wolfe; 1995:262–263.
 27. Castle WM, North PM. *Statistics in Small Doses*, 3rd ed. Edinburgh, England: Churchill Livingstone; 1995;17:60–61.