

المراجع

The Reviews

المراجع العربية:

القرآن الكريم، سورة الإنسان، الآية: 17.

القرآن الكريم، سورة المرسلات، آية: 20-23.

ابن القيم الجوزية (1987) زاد المعاد في هدي خير العباد - الجزء: 14 - الطبعة: 14، مؤسسة الرسالة - بيروت.

أبو عيانة، رمزي عبد الرحيم (2004) أهمية الزنجبيل في الطب الحديث. مجلة الخفجي. مدينة الملك عبد العزيز للعلوم والتقنية. العدد الثاني..، ص: 36-37.

البار، محمد علي (1991) الجنين المشوه والأمراض الوراثية الأسباب و العلامات والأحكام. دار القلم. دمشق. دار المنارة. جدة. الطبعة الأولى..، ص: 138-142.

جابر، فاطمة عياش حسن (2009) دراسة نمو الكلية والخصية في الجرذان البيضاء تحت تأثير المضاد الحيوي ماكسيبيم (سيفيبيم هيدروكلوريد) قبل وبعد الولادة. رسالة ماجستير. كلية التربية للبنات. الأقسام العلمية. جامعة الملك عبد العزيز. جدة.

الجوهري، محمد الجوهرى محمود (1994) علم الأنسجة وتقنياته. مكتبة الملك فهد الوطنية. الطبعة الأولى..، ص: 267-305.

خليفة، سناء (2008) تأثير نبات الزنجبيل ولبن الإبل على التغيرات المستحبثة بعقار الاسيتريتين (النيوتريجاستون) في طحال الجرذان البيضاء. مقبول للنشر. مجلة الجمعية المصرية الألمانية لعلم الحيوان.

الدويك، جميل القدسي (2004) الزنجبيل شراب ودواء ووقاية وشفاء - الطبعة الأولى- فهرسة مكتبة الملك فهد الوطنية أثناء النشر.

الشريف، منى محمد (2010) التأثير العلاجي للزنجبيل على السمية الوراثية المحدثة بعقار التاكسول (مضاد للسرطان) في نخاع عظم ذكور الفئران. السجل العلمي. اللقاء الخامس والعشرون للجمعية السعودية لعلوم الحياة. جامعة الملك فيصل. الإحساء.

العراني، فاطمة سعد سالم (2007) تأثير نبات الزنجبيل ولبن الإبل على بعض أعضاء ذكور الجرذان البيضاء المعاملة بعقار الأسيتريتين لإظهار الإعجاز العلمي في السنة (دراسة نسيجية وكيمياء نسيجية). رسالة ماجستير. كلية التربية للبنات. الأقسام العلمية. جامعة الملك عبد العزيز. جدة.

كريم، صالح عبد العزيز (1990) المدخل إلى علم الأجنة الوصفي والتجريبي. دار المجتمع للنشر والتوزيع. جدة. الطبعة الأولى.

كريم، صالح عبد العزيز والقدسى، فاطمة محمد (2008) علم الأجنة الوصفي المقارن. مركز النشر العلمي. جامعة الملك عبد العزيز. جدة. الطبعة الأولى.

لبنية ، محيي الدين (2005) الزنجبيل تابل ودواء. مجلة الخفجي. مدينة الملك عبد العزيز للعلوم والتكنولوجيا. العدد العاشر. ص:30-33.

نصر، أحمد نعمان (1995) علم الأنسجة. دار المعارف- القاهرة. الطبعة الثانية.

هرسانى، أحلام عباس محمود (2010) تأثير عقار الكربامازيبين (الترجيتول) على نمو الكبد والمبيض في الجرذان البيضاء. رسالة ماجستير. كلية العلوم للبنات. جامعة الملك عبد العزيز. جدة.

اليوسف، ز. و. (2002) دراسة الشكل الظاهري والنسيجي لتأثير عقار السيفوتاكسيم وعقار التراميسين على نمو المخ المتوسط في أجنة الدجاج. رسالة دكتوراه. كلية التربية للبنات- وكالة الرئاسة للكليات البنات . الرياض

المراجع الأجنبية:

A

Abd El Fatah ,A.A.(1992) Effect of Tetracyclines On the development of the liver in chick embryos. Ph .D.Theesis ,Al Azhar university, Cairo.Egypt.

Abdel-Aziz, H., Nahrstedt, A., Peterelt, F., Windeck, T., Ploch, M and Verspohl, E.J. (2005) 5-HT3 receptor blocking activity of arhyalk-anes isolated from the rhizome of *Zingiber officinale*. Planta. Med; 71: 609 – 616 .

Abebe W. (2002) Herbal medication: potential for adverse interactions with analgesic drugs. J Clin Pharm Ther. 27(6):391-401.

Aeschbach, R., Loliger, J., Scott, BC., Murcia, A., Butler, J., alliwell, B and Aruoma, OI. (1994) Antioxidant actions of Thymol, Carvacrol, 6-Gingerol, Zingerone and Hydroxythyrosol. Fd Chem. Toxic; 32(1): 31 - 36 .

Ahmed, R.S. and Sharma, S.B. (1997) Biochemical studies on combined effects of garlic (*Allium sativum Linn*) and ginger (*Zingiber officinale Rosc*) in albino rats. Indian J Exp Biol. 35, pp. 841–843.

Ahmed, RS., Seth, V and Banerjee, BD. (2000) Influence of dietary ginger system in rat: Comparison with ascorbic acid. Indian Journal of Experimental Biology; 38: 604 – 606.

Ahmed,B.F. , Dawood,A.A.,AbouAywa,E,H. and Ahmed, N.F.(1994) (Antenatal effect Of piroxicam(feldene) on kidney of white rat.Sc.J.A2.Med.Fac.(Girls). Vol. 15.No.1:569-579.

Ajith TA, Aswathy MS, Hema U. (2008) Protective effect of *Zingiber officinale* roscoe against anticancer drug doxorubicin-induced acute nephrotoxicity. Food Chem Toxicol. 46(9):3178-81.

Ajith, T.A., V. Nivitha and S. Usha, (2007) *Zingiber officinale* Roscoe alone and in combination with alpha-tocopherol protect the kidney against cisplatin-induced acute renal failure. Food. Chem. Toxicol., 45(6): 921-927.

Akil ,I.;Inan ,S.;Gurcu,B.; Nazikoglu,A.and Ozbilgin,K. (2004) (Histopathological and Ultrastructural Effects of losartan on embryonic rat kidney). Acta histochemica .107.291-300.

Alexander, A. F. et al. (1984) Oxsytetracycline associated nephrotoxicosis in feedlot calves. J. Am. Vit. Mid. Assoc., 185, 793-795.

Ali, B.H., Blunden, G., Tanira, M.O. and Nemmar, A. (2008) Some phytochemical, pharmacological and toxicological properties of ginger(*Zingiber officinale* Roscoe): A review of recent research. Food and Chemical Toxicology. Volume 46, Issue 2,P:409-420.

Al-Qattan, K., Thomson, M., and Alib, M. (2008) Garlic (*Allium sativum*) and ginger (*Zingiber officinale*) attenuate structural nephropathy progression in streptozotocin-induced diabetic rats. e-SPEN, the European e-Journal of Clinical Nutrition and Metabolism. Volume 3, Issue 2, P:e62-e71.

Al-Yahya, MA. Rafatulla, S. and Mossa, JS. (1989). Gastro Protective Activity of Ginger on Albino Rats, Am. J. Chi. Med. 17: 51-56.

Al-Zubairi AS, Abdul AB, Syam MM. (2010) Evaluation of the genotoxicity of zerumbone in cultured human peripheral blood lymphocytes. Toxicol In Vitro. 24(3):707-12.

Amin, A. and A.A. Hamza, (2006) Effects of Rosell and ginger on cisplatin- induced reproductive toxicity in rats. Asian J. Androl., 8(5): 607-612.

Amin,R.; Vickers,A.; Sistare,F.; Thompson,L.; Roman,J.; Lawton,M.; Kramer,J; Hamadeh,K.; Collins,J.; Grissom, S.;Bennett,L. ;Tucker,J. and Tennant,R. (2004) (Identification of putative Gene – based Markers of renal Toxicity). Environmental Health perspectives.Volume112,465-479 .

Ansari, M.N., U. Bhandari and K.K. Pillai, (2006) Ethanolic *zingiber officinale* R. extract pretreatment alleviates isoproterenol –induced oxidative myocardial necrosis in rats. Indian J. Exp. Biol., 44(11): 892-897.

Anufrieva ,R.G.;Zel tser ,I.Z.;Balabanova,E.L.; lapchinskaia A.V.;Baru,R.v.(1980) (Experimental study of the effect of rifamicin on the embryogenesis of White rats). Antibiotiki 25(4):280-4.

Ariens,E.J.;Simons,A.M.;and Offermeier,J.(1976) (Introduction to General Toxicity) Academic Press, Inc. New York.

Assael,B.M.(1987) (Toxic side effects of antibiotics in neonate:problems of oto and Nephrotoxicity of the aminoglycosides.V. Mediteranean congress of Chemotherapy (26Oct-1Nov;1986).cairo.Egypt Edited by Sabbour M.S and Cocuzza G., Supplemeut No2vol.6:336.

Aupee O, Almeras D, Le Garlantezec P, Bohand X. (2009) [Doxycycline]. Med Trop (Mars). 69(6):556-8.

Awodele O, Akintonwa A, Osunkalu VO, Coker HA. (2010) Modulatory activity of antioxidants against the toxicity of Rifampicin in vivo. Rev Inst Med Trop Sao Paulo. 52(1):43-6.

Azzam EI, Yang Z, Li M, Kim S, Kovalenko OA, Khorshidi M, Ende N. (2010) The effect of human cord blood therapy on the intestinal tract of lethally irradiated mice: possible use for mass casualties. Int J Radiat Biol.86(6):467-75.

B

Backon, J. (1991) Ginger as an antiemetic: Possible side effects due to its throm-boxane synthetase activity. Anaesthesia; 46: 705 – 706.

Balch, J.F. and Balch, P.A. (1996) Prescription for nutritional healing: A practical A-Z reference to drug-free remedies using vitamins, minerals, herbs & food supplements, Avery Publishing Group, New York.

Balubid, S. O. (2006) Effect of camel's urine and milk at the time of gestation and lactation on growth of kidney and liver on the rats. arabiologists. 25: 38-49.

Bankston, P.W. and Pinom, R.M.(1980) The development of the sinusoids of fetal rat liver: morphology of endothelial cells, Kupffer cells, and the transmural migration of blood cells into the sinusoids. Am J Anat. 159(1):1-15.

Barinov,E.F. and Sulaeva,O.N.(2003) (Morphometric characteristics of the kidney Nephrogenic zone in newborn rats) Morfologija. 123(2):77-80.

Barnes, J. (2003) Complementary therapies in pregnancy. The Pharmaceutical. J., (6):402-404.

Barth RK, Gross KW, Gremke LC, Hastie ND.(1982) Developmentally regulated mRNAs in mouse liver. Proc Natl Acad Sci U S A.79(2):500-4.

Beck,H.; Nierhoff,N.; Sack,K. (1980) (Experimental studies in animals on the nephrotoxicity of some new cephalosporin antibiotics: cefamandol, EMD29 645 and 29 946 (author ,s transl) Arzeimittelforschung. 30(4):660-6.

Beckstrom-Sternberg D. J. (1999) Dr. Duke's Phytochemical, and Ethnobotanical Databases. Phytochemical Database, USDA-ARS-NGRL, Beltsville Agricultural Research Centre (<http://www.ars-grin.gov/cgi-bin/duke>).

Berezhinskaia,V.V.;Dolgova,G.V.;Egorenko,G.G.;Svino ge eva,T.P. ;Zebrev,A.I.;Smolkina,T.V.; Shtegelman,L.A. and Nikitin,A.V. (1990) (General toxic and organotropic properties of cefotaxime in acute and chronic experiments)Antibiot khimoter.35(5):25-7.

Berman B, Perez OA, Zell D. (2007) Update on rosacea and anti-inflammatory-dose doxycycline. Drugs Today (Barc). 43(1):27-34.

Berndt,W.O.(1989) (Potential involvement of renal transport mechanisms in nephro toxicity) Toxicology Letters, Volume 46,lssues1-3p77-82.

Betz O, Kranke P, Geldner G, Wulf H, Eberhart LH. (2005) [Is ginger a clinically relevant antiemetic? A systematic review of randomized controlled trials]. Forsch Komplementarmed Klass Naturheilkd. 12(1):14-23.

Bhandari, U., J.N. Sharma and R. Zafar, (1998). The protective action of ethanolic ginger (*zingiber officinale*) extract in cholesterol fed rabbits. J. Ethanopharmacol., 61(2): 167-171.

Black CD, Herring MP, Hurley DJ, O'Connor PJ. (2010) Ginger (*Zingiber officinale*) Reduces Muscle Pain Caused by Eccentric Exercise. J Pain. [Epub ahead of print].

Blanchard P, Rudhardt M, Fabre J, Leski M. (1977) [Pharmacokinetic effect of various antibiotics in the tissues]. Schweiz Med Wochenschr. 107(17):589-97.

Blumenthal M, Busse W, Goldberg A, et al, eds. (1998) The Complete German Commission E Monographs: Therapeutic Guide to Herbal Medicines. Austin, TX: American Botanical Council.

Bode, AM., Ma, WY., Surch, YJ and Dong, Z. (2001) Inhibition of epidermal growth factor induced cell transformation and activator protein 1 activation 6-gingerol. Cancer Res., 1 ; 61 (5): 850 - 3 .

Bokor-Bratić M, Brkanić T. (2000) [Clinical use of tetracyclines in the treatment of periodontal diseases]. Med Pregl. 53(5-6):266-71.

Bolender, R. P. and Weibel, E. R. (1973) A morphometric study of the removal of phenobarbital-induced membranes from hepatocytes after cessation of treatment. J Cell Biol. 56:746.

Bone, ME., Wilkinson, DJ., Young, JR and McNeil, J. (1990) Ginger root- a new antiemetic: the effect of ginger root on postoperative nausea and vomiting after major gynecological surgery. Anaesthesia; 45: 669 – 71.

Bordia A, Verma SK, Srivastava KC. (1997) Effect of ginger (*Zingiber officinale* Rosc.) and fenugreek (*Trigonella foenumgraecum* L.) on blood lipids, blood sugar and platelet aggregation in patients with coronary artery disease. Prostaglandins Leukot Essent Fatty Acids. 56(5):379-84.

Borlak,J.(2005) (Handbook of Toxicogenomics: Strategies and Applications (chapter 19).wiley –VCH Verag GmbH and Co KkaA,weinheim, pp:471-486.

Borrelli F, Capasso R, Aviello G, Pittler MH, Izzo AA. (2005)
Effectiveness and safety of ginger in the treatment of pregnancy-induced nausea and vomiting. Obstet Gynecol. 105(4):849-56.

Breen KJ, Schenker S, Heimberg M. (1975) Fatty liver induced by tetracycline in the rat. Dose-response relationships and effect of sex. Gastroenterology. 69(3):714-23.

Bergner, P. (1991) Is ginger safe in pregnancy?. Med Herbalism. 3:3.

Brent,R.L. and Jersh,R.P .(1976) (Intrauterine growth Retardation Adv. Teratology, 2:139-227.

C

Centers for Disease Control and Prevention (CDC). (2001) Update: Interim recommendations for antimicrobial prophylaxis for children and breastfeeding mothers and treatment of children with anthrax. MMWR Morb Mortal Wkly Rep. 50(45):1014-6.

Chaibi, A. (1984) [Administration of tetracycline to the intact and thyroidectomized rat: its effect on the ultrastructure of hepatocytes, lipids in the liver and plasma]. Arch Inst Pasteur Tunis. 61(1):51-72.

Chambers,H.F.(2001) (Antimicrobial Agents,General consideration) Chapter 43 in (Goodman and Gilmams Basis of therapeutics) 10th Edition Mc Graw - Hill, Medical Publishing division New York, London Toronto.

Cheville, N. F. (1983) Cell pathology. Second ed. Chapter 9. The Iowa State University Press, Ames, Iowa, USA.

Chrubasik, S.; Pittler, M.H. and Roufogalis, B.D., (2005) Zingiberis rhizome: a comprehensive review on the ginger effect and effecact profiles. Phytomedicine. 12(9):684-701.

Chung, WY., Yow, CM and Benzi, IF. (2003) Assessment of membrane protection by traditional Chinese medicine using aflow cytometric technique: preliminary findings. Redox. Rep. 8 (1): 31- 3.

Cohlan.S.Q. et al. (1963) Growth inhibition of prematures receiving tetracycline. Am. J. Dis. Child.,105:65-73.

Colalto C. (2010) Herbal interactions on absorption of drugs: Mechanisms of action and clinical risk assessment. Pharmacol Res.

Convington,T.R.;Dipalma,J.R.;Hussrv,D.A.;Lasagna,L;Latro, D.S.and whitsett,T.L. (1985) (Drug facts and comparisons). A divition of J B Lippinott company phiadelphia Toronto ,P:1350.

Criag, WJ. (1997) Phytochemical: guardians of our health. J. Am. Diet. Assoc; 97 (10 supp 12): 199 - 204 .

Culig J, Mlinarić-Dzepina A, Leppée M, Vranes J. (2010) [Resistance of uropathogenic strains of Escherichia coli in pregnant women and other women in generative ages in comparison with antibiotics consumption in Zagreb]. Med Glas Ljek komore Zenicko-doboj kantona. 7(1):54-9.

Curran,R.C.(1996) Coloun atlas of histopathology . 3rd ed .Oxford university press. New york. Pp280.

Czeizel AE, Rockenbauer M. (2000) A population-based case-control teratologic study of oral oxytetracycline treatment during pregnancy. Eur J Obstet Gynecol Reprod Biol. 88(1):27-33.

Czeizel AE, Rockenbauer M. (1997) Teratogenic study of doxycycline. Obstet Gynecol. 89(4):524-8.

D

Dallner G, Siekevitz P, Palade GE.(1966) Biogenesis of endoplasmic reticulum membranes. II. Synthesis of constitutive microsomal enzymes in developing rat hepatocyte. J Cell Biol. 1966 Jul;30(1):97-117.

Damjanov, I. and Solter, D. (1971) Ultrastructure of acute tetracycline induced liver change .Experientia., 27:1204-1208.

Dashe, JS. and Gilstr, LC. 3rd. (1997) Antibiotic use in pregnancy. Obstet Gynecol Clin North Am. 24(3):617-29.

de Louvois J, James J, Mulhall A. (1984) Latamoxef and the newborn. Arch Dis Child. 59(4):346-50.

Demirkaya, E.; Atay, A.A.; Musabak, U.; Sengul, A. and Gok,F. (2006) (Ceftriaxone-related Hemolysis and acute renal failure). Pediatr Nephrol. 21(5):733-6.

Dickinson, H.; Walker,D.; Wintour,M. and Moritz,K. (2007) (Maternal dexamethasone treatment at midgesation reduces nephron number and alters renal gene expression in the fetal spiny mouse). American physiological Society 453-461.

Dickinson,H.;Walker,D.W.;Cullen-Mcewen,L.A. and Wintour,E.M. (2005) (the Spiny mouse(Acomys cahirinus) completes nephrogenesis before to birth). Am J physiol Renal physio. 1289:F273-F279.

DiGiacomo, J.E. and Hay, W.W. Jr, (1990) Fetal glucose metabolism and oxygen consumption during sustained hypoglycemia. Metab Clin Exper. 39, pp. 193–202.

Dog, T. L. (2009) the use of botanicals during pregnancy and lactation. Alternative Therapies, Vol. 15, No. 1.

Dugasani S, Pichika MR, Nadarajah VD, Balijepalli MK, Tandra S, Korlakunta JN. (2010) Comparative antioxidant and anti-inflammatory effects of [6]-gingerol, [8]-gingerol, [10]-gingerol and [6]-shogaol. J Ethnopharmacol. 127(2):515-20.

E

Edwards RK. (2000) Syphilis in women. Prim Care Update Ob Gyns. 7(5):186-191.

Edwards, B. R.; Mendel, D. B.; Larochelle, F. T. and Sternand Valtin, H. (1981) Postnatal development of urinary concentrating ability in rats: changes in renal anatomy and neurohypophysial hormones in Spitzer A, ed. The Kidney During Development Morphology and Function. New York, Masson, 233-240.

Egwurugwu, J. N., Ufearo, C. S., Abanobi, O. C., Nwokocha, C. R., Duruibe, J. O., Adeleye, G. S., Ebunlomo, A. O., Adetola, A. O. and Onwufaji, O. (2007) Effects of ginger (*Zingiber officinale*) on cadmium toxicity. African Journal of Biotechnology. Vol. 6 (18), pp. 2078-2082.

Ekwall B, Acosta D. (1982) In vitro comparative toxicity of selected drugs and chemicals in HeLa cells, Chang liver cells, and rat hepatocytes. Drug Chem Toxicol. 5(3):219-31.

EL-Ghawaby ,F.A.; Abd El-Wahab ,M.G.; El- Nabarawy,S.E. and Sadek ,S.A. (1994) (Effect of Cefotaxime on the developing kidney of the chick embryo) J.Biomed .Sci .Ther ;10:10-26.

El-Hummmdi, L.A.T. (2006) Therapeutic Effect of Honey with *Nigella Sativa* Mixure , and Ginger on The Toxic Impacts of Enoxaparin Sodium (Anticoagulant) On The Kidney Tissue of Albino Rats. J. Egypt. Soc. Toxicol. Special Issue. Su ppl., 34:130-137.

Ernst, E. (2002) The risk–benefit profile of commonly used herbal therapies: Gingko, St. John’s wort, ginseng, echinacea, saw palmetto and kava. Ann Int Med.136(1):42–53.

Ezeuko V. C. ; Nwokocha C. R. ; Mounmbegna P. E. and Nriagu C. C.(2007) Effects of *zingiber officinale* on liver function of mercuric chloride-induced hepatotoxicity in adult wistar rats. Electron J Biomed .3:40-45.

F

Fanning ,W.L.; Gump,D.; Jick, H.(1976) Gentamicin and cephalothin associated rises in blood urea nitrogen. Antimicrob Agents Chemother. 10:80-82.

Farid M. (1981) Alterations induced by Nesdonal and Revrin on the hepatic and nervous tissues of chick embryo. Pediatr Pharmacol (New York). 1(4):357-64.

Faridc,M.(1981) (Alterations induced by nesdonal and Revrin on the hepatic and nervous tissues of chick embryo). Pediatr.Pharmacol New York,Vol.1(4),P:357-64.

Ferjani, A.; Ben Abdallah, H.; Ben Saida, N.; Gozzi, C. and Boukadida, J., (2006) Viginal colonization of the Streptococcus agalactiae in pregnant woman in Tunisia: risk factors and susceptibility of isolates to antibiotics. Bull Soc Pathol Exot., 99(2):99-102.

Fischer-Rasmussen,W. Kjaer, S. Dahl, C. and Asping,U. (1990) Ginger treatment of hyperemesis gravidarum. Eur J Obstet Gynaecol Reprod Biol.38: 19–24.

Fogiel,M.(1989) The statistics proplom solver ,res .New jersy.

Forster DA, Denning A, Wills G, Bolger M, McCarthy E. (2006) Herbal medicine use during pregnancy in a group of Australian women. BMC Pregnancy Childbirth.6:21.

Friedman, JM. Little BB, Brent RL, Cordero JF, Hanson JW, Shepard TH. (1990) Potential human teratogenicity of frequently prescribed drugs. Obstet Gynecol. 75(4):594-9.

Friedman, J.M. and Polifka, J.E. (2000) (Teratogenic Effects of Drugs) A Source for Clinicians (TERIS) :Baltimore.MD:The John Hopkins university press.

Fuentes VO, Gonzalez H, Sanchez V, Fuentes P, Rosiles R. (1997) The effect of neomycin on the kidney function of the horse. Zentralbl Veterinarmed A. 44(4):201-5.

Fujita, K. and Fujita, HM. (1985) Reversibility of antibiotic incorporation in large organelles of kidney cell. Int J Clin Pharmacol Ther Toxicol. 23(6):291-4.

Fulder S, Tenne M. (1996) Ginger as an anti-nausea remedy in pregnancy: the issue of safety. Herbalgram. 38: 47–50.

G

Ganapathy ,M.E.;Huang,W.and Rajan D.P.(2000) (Beta-lactam antibiotics as substrates for OCTN2 ,an organic cation/carnitine transporter). J Biol Chem. 275:1707.

Gartner,L.P.and Hiatt,J.L.(2006) (Color Atlas of Histology). 4th ed. Lippincott Williams and Wilkins. Newyork .London.

Geall, M. G. and Webb, M. J. (1974) Liver disease in pregnancy. Medical Clinics of North America. Vol. 58, No. 4:817-822.

Ghadially, F. N. (1988). Ultrastructural pathology of the cell and matrix. A text and atlas of physiological and pathological alterations in the fine structure of cellular and extracellular components. 3rd ed. Volume I. Butterworth's.

Ghayur MN, Khan AH, Gilani AH. (2007) Ginger facilitates cholinergic activity possibly due to blockade of muscarinic autoreceptors in rat stomach fundus. Pak J Pharm Sci. 20(3):231-5.

Gilbert,T.;LeLiever-Pegorier M.and Merlet- Benichou,C. (1990) (Immediat and Long-term renal effects of foetal exposure to gentamicin.J.Pediat-Nephrol, 4(4):445-450.

Goodman and Gilmans (2001) The Pharmacological Basis of Therapeutics 10 Edition. Joel, G.; Hardman L.E. E Limbird New York, London.P:1240-1246.

Goyal, RK and Kadnur, SV. (2006) Beneficial effects of Zingiber officinale on goldthioglucose induced obesity. Fitoterapia. 77(3):160-3.

Gray, I. E., Weaver, R. N., Skinner, P. Mathews, J., Day, C.E. and Stern, K. (1974) Effects of tetracycline on Ultrastructure and lipoprotein secretion in the rat hepatocytes. Taxicol. Appl. Pharmacol. 317-332.

Green DM, Norkool P, Breslow NE, Finklestein JZ, D'Angio GJ. (1990) Severe hepatic toxicity after treatment with vincristine and dactinomycin using single-dose or divided-dose schedules: a report from the National Wilms' Tumor Study. J Clin Oncol. 8(9):1525-30.

Gregorio GV, Ball CS, Mowat AP, Mieli-Vergani G. (1993) Effect of rifampicin in the treatment of pruritus in hepatic cholestasis. Arch Dis Child. 69(1):141-3.

Guahk GH, Ha SK, Jung HS, Kang C, Kim CH, Kim YB, Kim SY. (2010) Zingiber officinale Protects HaCaT cells and C57BL/6 Mice from Ultraviolet B-Induced Inflammation. J Med Food. 13(3):673-80.

Guggenbichler JP, Schabel F. (1979) [Tetracyclin intoxication versus idiopathic pancreatitis: report of a case with multiple organ involvement (author's transl)]. Padiatr Padol. 14(3):267-72.

Guglielmotti MB, Ubios AM, Larumbe J, Cabrini RL. (1989) Tetracycline in uranyl nitrate intoxication: its action on renal damage and U retention in bone. Health Phys. 57(3):403-5.

Juliaev AE, Kivman GIa, Nadirova BA. (1982) [Role of age in the interaction of tetracycline with the subcellular fractions of rat liver homogenate]. Antibiotiki. 27(10):776-8.

Guzmán A, García C, Marín AP, Willoughby C, Demestre I. (2003) Developmental toxicity studies of the quinolone antibacterial agent irloxacin in rats and rabbits. Arzneimittelforschung. 53(2):121-5.

Gwee MC. (1982) Can tetracycline-induced fatty liver in pregnancy be attributed to choline deficiency?. Med Hypotheses. 9(2):157-62.

H

Habib,S. H. M. ; Makpol, S. ; Abdul Hamid, N. A. ; Das,S. ; Ngah, W. Z. W. and Yusof, Y. A. M. (2008) Ginger Extract (Zingiber Officinale) has Anti-Cancer and Anti-Inflammatory Effects on Ethionine-Induced Hepatoma Rats. Clinics. 63(6): 807–813.

Hagel-Lewicka E, Lewicki Z, Wutzen J, Figurski R. (1980) Ultrastructural investigations of the liver in young (suckling) rats receiving tetracycline hydrochloride. Mater Med Pol. 12(4):259-62.

Haksar A, Sharma A, Chawla R, Kumar R, Arora R, Singh S, Prasad J, Gupta M, Tripathi RP, Arora MP, Islam F, Sharma RK. (2006) Zingiber officinale exhibits behavioral radioprotection against radiation-induced CTA in a gender-specific manner. *Pharmacol Biochem Behav.* 84(2):179-88.

Halliwell, B. (1996) Antioxidant in human health and disease. *Annu. Rev. Nut.,* 16: 33-38.

Hamilton, W.J. and Massman, H.W. (1972) *Human Embryology. Prenatal development of form and function.* Fourth Edition. Published by the Macmillan Press LTD The Williams and Wilkins Company, P: 340-362.

Han, S.S. and Holmsted, J.V. (1981) *Human Microscopic Anatomy.* First Edition. Published by McGraw-Hill Book Company New York St. Lows San Francisco, Auckland, Bongota, Harmburg, Johannesburg, London, Madrid, Mexico, Montreal, New Delhi, Panama, Paris, Sao Paulo, Singapore, Sydeney, Tokyo, Toronto. P:449-461.

Hannah R, Sahib MK. (1975) Stabilization of rat liver tyrosine aminotransferase by tetracycline. *Biochem J.* 150(3):329-33.

Hara,K.;Muraoka,Y.;Yoshida,T.;Muranaka,R.;Kanamori, S.; Hirashiba,M; uchida,H. and Ikeuchi, K.(2001) (Reproductive and developmental toxicity studies of S-1090, Cefmatilen hydrochloride hydrate(1) –A study On oral administration prior to and in the early stages of pregnancy in rats) *J Toxicol Sci.*26 suppl 1:157-70.

Haschek,W.M. and Rousseaux,C.G.(1991) (*Urinary system In Handbook of Toxicologic Pathology*) Academic press, San Diego, California ,pp.315- 387(chapter15).

Hayashi, K. (1961) Experimental studies on histopathological changes in the liver and kidney of rats following antibiotics administration. *Tokushima J Exp Med.* 8:156-62.

Heeba GH, Abd-Elghany MI. (2010) Effect of combined administration of ginger (Zingiber officinale Roscoe) and atorvastatin on the liver of rats. *Phytomedicine.*

Hopf,G.; Bocker,B.; Holzmann,I.G. and Estler,C.J. (1985) Quantitative assessment of the polysome profile of the livers of mice treated with tetracycline or doxycycline. Bio.che.Pharm;34(20):3689-3701.

Hull,R.L. and Brandon ,D.(1991) Thrombo cytopenia possibly caused by structurally related third – generation cephalosporins. Dicp.25(2): 135-6.

Hutterer, F. et al. (1969) Hepatocellular adaptation and injury. Lab Invest 20:455.

Hyams, D. E. and Isselbacher, K. J. (1964) Presentation of fatty liver by administration of adenosine triphosphate. Nature. 204:1196-1964.

I

Idu SR, Zimmerman C, Mulder L, Meis JF. (1998) [A very serious course of psittacosis in pregnancy]. Ned Tijdschr Geneesk.d. 142(47):2586-9.

Infante, R., Koumanov, K. and Caroli, J. (1971) Tetracycline-induced fatty liver. Biochemical Mechanisms. Gut. 12:765-766.

Ippoushi K, Ito H, Horie H, Azuma K. (2005) Mechanism of inhibition of peroxynitrite-induced oxidation and nitration by [6]-gingerol. Planta Med. 71(6):563-6.

J

Jagetia, GC., Baliga, MS and Ulloor, JN. (2003) Influence of Ginger Rhizome (*Zingiber officinal Rosc*) on survival, glutathione and lipid peroxidation in mice after whole-body exposure to Gamma Radiation. Radiation Research. New York; 160: 584 - 592.

Jagetia G, Baliga M, Venkatesh P. (2004) Ginger (*Zingiber officinale Rosc.*), a dietary supplement, protects mice against radiation-

induced lethality: mechanism of action. Cancer Biother Radiopharm. 19(4):422-35.

Jagt CT, Zuckermann M, Ten Kate F, Taminiau JA, Dijkgraaf MG, Heij H, De Kraker J, Verschuur AC. (2009) Veno-occlusive disease as a complication of preoperative chemotherapy for Wilms tumor: A clinico-pathological analysis. Pediatr Blood Cancer. 53(7):1211-5.

Jaton, K. and Greub, G. (2005) [Chlamydia: diagnostic and treatment]. Rev Med Suisse ,1(13): 895-8,901-3.

Jayasagar,G.; kumar,K.; Rao,M.; Chandasekhar, K.; and Rao,M. (2002) (Effect of Cephalexin on the pharmacokinetics of metformin In healthy human volunteers). Drug metabol Drug Interact .19(1):41-8.

Journey, LJ. and Goldstein, MN. (1963) The effect of terramycin on the fine structure of HeLa cell mitochondria. Cancer Res. 23:551-4.

Junqueira, L. C. ; Carneior, J. and Long, J. A.(1998) Basic Histology. 10th ed.

K

Kacet N, Zaoui C, Roussel-Delvallez M, Dubos JP, Martin GR, Lequien P. (1986) [Piperacillin in the newborn infant. A clinical and pharmacologic study]. Presse Med. 15(46):2339-41.

Kai,S.;Kohmura,H.;Kawano ,S.;Sakai,A.;Kuroyanagi, K. and Kadota ,T.(1992) (Reproductive and Developmental toxicity studies on cefepime Dihydrochloride administered subcutaneously to Rats during the premating gestation and lactation Periods) J Antibiot. 45(6): 642-60.

Kanamura, S.; Kanai, K. and Watanabe, J. (1990) Fine structure and function of hepatocytes during development. J. Electron Microsc. Tech., 14: 92.

Kanerva, L., Estlander, T and Jolanki, R. (1996) Occupational allergic contact dermatitis from spices. Contact Dermatitis; 35: 157 – 162.

Kasabulatov,K.K.;manuilov,K.K.and Voropaeva, S.D.(1993) (An animal experiment On transplacental transport of Ceftazidim at various) .Antibiotc khimioter. 3891:33-8.

Kato,A. ; Higuchi, Y. ; Goto, H. ; kizu,H. ; Okamoto,T. ; Asano,N. ; Hollinshead,J. ; Nash,R. J. and Adachi, I. (2006) Inhibitory Effects of Zingiber officinale Roscoe Derived Components on Aldose Reductase Activity in Vitro and in Vivo. J. Agric. Food Chem. 54:664-□6644.

Kelly, K.J.; Sutton, T.A.; Weathered, N.; Ray, N.; Caldwell, E.J.; Plotkin, Z. and Dagher, P.C. (2004) Minocycline inhibits apoptosis and inflammation in a rat model of ischemic renal injury. Am. J. Physiol Renal Physiol., 287(4): F760-6.

Khalifa, S. A. (2006) Effect of camel's Urine and milk, Honey Bee with Nigella Sativa Mixture and Ginger on Toxic Potentials of Haloperidol (Antipsychotic Agents) on Fertility in the Male Albino Rats. J. Egypt. Soc. Toxicol, Special Issue.,34:119-12.

Khalifa, S. A. (2007) Protective effects of the ginger and the honey bee with Nigella Sativa on the clexane - induced pathological changes in duodenum of rats. Egypt. J. Comp. Path. & Clinic. Path. Vol; 20(1): 81 - 91 .

Khatib R, Thirumoorthi MC, Kelly B, Grady KJ. (1995) Severe psittacosis during pregnancy and suppression of antibody response with early therapy. Scand J Infect Dis. 27(5):519-21.

Kierszenbaum, A.L.(2002) Histology and cell biology: An introduction to pathology. Mosby, Inc. USA.

Kirchner, G. ; Habers, M. ; Bunsch, A. ; Seitz, H.J. and Hoppner, W. (1993) Zonation of glucokinase in rat liver change during postnatal development. Febs Lett, 328(1-2): 119-124.

Kirjuschenkov AP. (1978) [Effect of antibiotics on the early stages of embryogenesis]. Zentralbl Gynakol. 100(22):1451-3.

Kleinschmidt S, Rump G, Kotter J. (2007) [Herbal medications. Possible importance for anaesthesia and intensive care medicine] Anaesthetist. 56(12):1257-66.

Knothe, H. Dette , G. A. (1985) Antibiotics in Pregnancy: Toxicity and Teratogenicity. Infection. 13:3-49.

Koch C, Reichling J, Schneele J, Schnitzler P. (2008) Inhibitory effect of essential oils against herpes simplex virus type 2. Phytomedicine. 15(1-2):71-8.

Kohda,Y.and Gemba,M. (2005) (Cephaloridine induces translocation of protein kinase C into mitochondria and enhances mitochondrial generation of free radicals In the kidney cortex of rats causing renal dysfunction) J Pharmacol ;98(1):49-57.

Kondrackienė, J., Kupčinskas, L. (2008) Liver diseases unique to pregnancy. Medicina (Kaunas); 44(5):337-345.

Korzeniowski OM. (1995) Antibacterial agents in pregnancy. Infect Dis Clin North Am. 9(3):639-51.

Kota, N., Krishnaa, P. and Polasaa, K. (2008) Alterations in antioxidant status of rats following intake of ginger through diet. Food Chemistry. Volume 106, Issue 3, P:991-996.

Kovnat,P,Labovits,E. and Levison,S.P. (1973) Antibiotics and the kidney .Med. Clin.North Am.57:1045-1063.

Kraus RL, Pasieczny R, Lariosa-Willingham K, Turner MS, Jiang A, Trauger JW. (2005) Antioxidant properties of minocycline: neuroprotection in an oxidative stress assay and direct radical-scavenging activity. J Neurochem. 94(3):819-27.

Kumar,V.;Cotran,R.S. and Robbins,S.L. (2003) (Basic pathology). 7th Edition W.B.Saunders company. Australia ,Canada, UK/Europe, Japan or USA.

Kurebe,M.;Asaoka,H.;Hata,T.;Izaw,M.;Watanabe,T. and Sawazaki,S .(1984) (Toxicological studies on a new cephamycin,MT-141,X.Its prenatal and postnatal test in rats). J Antibiot. 37(6): 1211-27.

Kuwayaman,H. ;Nakajima,N.;Matsuo,Y. and Eastwood, G.L.(1990)
(Effects of single parenteral indomethacin injection in rat. Fundic and antral epithelial proliferation) J.Clin.Gastroenterol .12: 72-75.

L

Labbé G, Fromenty B, Freneaux E, Morzelle V, Letterson P, Berson A, et al. (1991). Effects of various tetracycline derivatives on in vitro and in vivo beta-oxidation of fatty acids, egress of triglycerides from the liver, accumulation of hepatic triglycerides, and mortality in mice. Biochem Pharmacol .41:638-641.

Langmead, L and Rampton, DS. (2001) Review article: herbal treatment in gastrointestinal and liver disease- -benefits and dangers. Aliment Pharmacol Ther; 15 (9): 1239 - 52.

Larson JL, Wallace TL, Tyl RW, Marr MC, Myers CB, Cossum PA. (2000) The reproductive and developmental toxicity of the antifungal drug Nyotran (liposomal nystatin) in rats and rabbits. Toxicol Sci. 53(2):421-9.

Lawrence ,D. (1978) (Immunolgic cross-Reactivity between penicillins and cephalosporins) .The Journal of Infectious Diseases .Vol ,137.PP.74-79.

Leung, A.Y. (1984) Chinese Herbal Remedies, Universe Books, New York.

Lewis, M.; Schenker, S. and Coombes, B., (1967) Studies on the pathogenesis of tetracycline induced fatty liver. Am. J. Dig. Dis., 12:429-435.

Liu, N., Huo, G., Zhang, L and Zhang, X. (2003) Effect of *Zingiber officinal* Rosc on lipid Peroxidation in hyperlipidemia rats. Wei Sheng Yan Jiu; 32 (1) : 22 - 3.

Loginov AV, Polosova RG, Poliak MS, Iukhlov AK. (1976) [Functional state of the adrenal cortex in the antibiotic therapy of experimental staphylococcal infection]. Antibiotiki. 21(7):631-6.

Lu P, Lai BS, Liang P, Chen ZT, Shun SQ. (2003) [Antioxidation activity and protective effect of ginger oil on DNA damage in vitro]. Zhongguo Zhong Yao Za Zhi. 28(9):873-5.

Luft,F.G.;Patel,V.;Yum,M.N.;Kleit,S.A. (1976) (Nephrotoxicity of cephalosporin gentamicin combinations in rats Antimicrob. Agents Chemother.9:831-839.

Luzzatto, A. C. (1981) Hepatocyte differentiation during early fetal development in the rat. Cell Tissue Res. 1981;215(1):133-42.

M

Machado, A. L. DS. ; Brando, A.A.H. ; da Silva, C.M.O. and da Rocha, R.F. (2003) Influence of tetracycline in the hepatic and renal development of rat's offspring. Brazilian Archives of Biology and Technology.Vol.64,n.1:pp.47-51.

MacKintosh D, Baird-Lambert J, Drage D, Buchanan N. (1985) Effects of prenatal glucocorticoids on renal maturation in newborn infants. Dev Pharmacol Ther. 8(2):107-14.

Mal'tsev GIu, Nesterin MF. (1978) [Effect of tetracycline on rat liver mitochondrial oxidative phosphorylation in vivo and in vitro under a varying body allowance of protein]. Vopr Pitan. (4):40-5.

Mallikarjuna K, Sahitya Chetan P, Sathyavelu Reddy K, Rajendra W. (2008) Ethanol toxicity: rehabilitation of hepatic antioxidant defense system with dietary ginger. Fitoterapia. 79(3):174-8.

Mark J. E. ; Thomas G. H. and Allen F. H. (2005) Effects of Antibiotics and Probiotics on Suckling Pig and Weaned Pig Performance. Intern J Appl Res Vet Med . Vol. 3, No. 4.

Marlier,A. and Gilbert,T.(2004) (Expression of retinoic acid-synthesizing and metabolizing enzymes during nephrogenesis in the rat). Gene Expr patterns. 5:179-185.

Matsuda A, Wang Z, Takahashi S, Tokuda T, Miura N, Hasegawa J. (2009) Upregulation of mRNA of retinoid binding protein and fatty acid binding protein by cholesterol enriched-diet and effect of ginger on lipid metabolism. Life Sci. 84(25-26):903-7.

Mayura K, Parker R, Berndt WO, Phillips TD. (1984) Effect of simultaneous prenatal exposure to ochratoxin A and citrinin in the rat. J Toxicol Environ Health. 13(4-6):553-61.

McMahon, C. MS and Pergament,E. (2001) The Treatment of Nausea and Vomiting in Pregnancy (NVP). Vol. 8, No. 2.

Mcreddie,A.;Romano,L.M.;Harris,J.M.; Ferder,L. and Gomez,R.A. (1995) (Angiotensin II regulates nephrogenesis and renal vascular development) American Physiological Society P110-115.

Mehrdad M, Messripour M, Ghobadipour M. (2007) The effect of ginger extract on blood urea nitrogen and creatinine in mice. Pak J Biol Sci. 10(17):2968-71.

Meijer A, Brandenburg A, de Vries J, Beentjes J, Roholl P, Dercksen D. (2004) Chlamydophila abortus infection in a pregnant woman associated with indirect contact with infected goats. Eur J Clin Microbiol Infect Dis. 23(6):487-90.

Meléndez Camargo ME, Lozano Cortés M. (1998) Renal and hepatic interactions of acetaminophen and amikacin in the infant rat. Proc West Pharmacol Soc. 41:61-3.

Meyer, K., Schwartz, J., Crater, D and Keyes, B. (1995) *Zingiber officinal* (ginger) used to prevent 8-Mop associated Nausea. Dermatology. Nursing; 7(4): 242 - 244.

Miller, S.E.P., Mac Sween, R.N.M., Glen, A.C.A. Tribedi, K. and Moore, E.M. (1967) Experimental studies on the hepatic effects of tetracycline. Brit. J. Exp. Path. 48:51-57.

Morrissey, R.E.; Tyl, R.W.; Price, C.J.; Ledoux, T.A.; Reel, J.R.; Paschke, L.L.; Marr, M.C. and Kimmel, C.A., (1986) The developmental toxicity of orally administered oxytetracycline in rats and mice. Fundam Appl Toxicol.,7(3):434-43.

Mosca A, Russo F, Miragliotta G. (2006) In vitro antimicrobial activity of benzalkonium chloride against clinical isolates of *Streptococcus agalactiae*. J Antimicrob Chemother. 57(3):566-8.

Mucklow, JC. (1986) The fate of drugs in pregnancy. Clin Obstet Gynaecol. 13(2):161-75.

Murata, P., Kase, Y., Ishige, A., Sasaki, H., Kurosawa, S and Nakamuvre, T. (2000) The herbal medicine Dai-kenchu-to and one its active components [6]-shogol increase intestinal blood flow in rats. Life Sci., 70 (17): 2061 - 70 .

Murphy, P.A. (1998) Alternative therapies for nausea and vomiting of pregnancy. Obstet Gynecol 91 pp. 149–155.

N

Nagai,J. andTakano,M.(2004) (Molecular aspects of renal handling of aminoglycosides And strategies for preventing the nephrotoxicity .Drug Metabolism and pharmacokinetics.19(3):159-170.

Nagasawa H, Watanabe K, Inatomi H. (2002) Effects of bitter melon (*Momordica charantia* L.) or ginger rhizome (*Zingiber officinale* rosc) on spontaneous mammary tumorigenesis in SHN mice. Am J Chin Med. 30(2-3):195-205.

Nakamura Y, Yoshida C, Murakami A, Ohigashi H, Osawa T, Uchida K. (2004) Zerumbone, a tropical ginger sesquiterpene, activates phase II drug metabolizing enzymes. FEBS Lett. 572(1-3):245-50.

Nakazawa T, Ohsawa K. (2002) Metabolism of [6]-gingerol in rats. Life Sci. 70(18):2165-75.

Nammi S, Kim MS, Gavande NS, Li GQ, Roufogalis BD. (2010) Regulation of Low-Density Lipoprotein Receptor and 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase Expression by *Zingiber officinale* in the Liver of High-Fat Diet-Fed Rats. Basic Clin Pharmacol Toxicol. 106(5):389-395.

Nardiello S, Pizzella T, Arivielo R. (2002) [Risks of antibacterial agents in pregnancy]. Infez Med. 10(1):8-15.

Nathanson,S.;Moreau,E.;Merletbenichou,C.and Gilber,T. (2000)
(In utero and in Vitro exposure to beta -lactams impair kidney development in the rat). J A M. 11(5):874-84.

Nelson, M.L. (1998) Chemical and biological dynamics of tetracyclines. Adv Dent Res 12:5-11.

Nelson, O. E. (1953) Comparative embryology of the vertebrates. 1st ed. Constable and company Limited 10 Orange. Street London, W.C.2:622-628.

Nie H, Meng LZ, Zhang H. (2006) [Effect of gingerol on endotoxemia mouse model induced by heatstroke]. Zhongguo Zhong Xi Yi Jie He Za Zhi. 26(6):529-32.

Nichoga,L.A.; Skosyрева,A.M.and Вово пеава,S.D.(1982)
Transplacental penetration Of gentamicin and its effect on foetus. J.Antibiotiki(Mosc). 27 (12):226-230.

Nigam, S. K.; Aperia, A. C. and Brenner, B. M. (1996)
Developmental and maturation of the kidney. In Brenner BM, ed
The kidney philadelohia, WB Saunders, 72-98.

Ninomiya,K. ;Haseyawa,Y. ;Kanamoto,T. ;Yoshimoto,Y.; Hamaya,K.;Masuda,K.and Nishio,Y.(1982) (Fundamental and clinical studies of cefotaxime during The perenatal period in pregnant women .JPN. Antibiotic ,35:1882-1892.

Nordeng, H. and Havnen, G.C. (2004) Use of herbal drugs in pregnancy: a survey among 400 Norwegian women. Pharmacoepidemiol Drug Saf., 13(6): 371-80.

Nomura A, Furuhashi T, Komuro E, Uehara M, Miyoshi K, Nakayoshi H. (1982) [Reproduction study on netilmicin. (1) Teratological study in rats (author's transl)]. Jpn J Antibiot. 35(2):614-29.

O

Olin,B.R.;Hebel,S.K.; Donbek,C.E.; Cremp,J.L. and Hulber, M.K.(1995) (Drug Facts and Comparisons) 49th Edition published by facts and comparisons.P. 2763-3201.

P

Page, C.P.; Cartis ,M.J.; Suyyer, M.C.; Walker, J. and Hoffman, M.B.B. (1997) (Mosby Integrated pharmacology) London .Chicago.Philadephia.

Paget,G.E. and Barnes,J.M. (1964) Evalution of Drug Activites, and pharmacokinetics. Academic press; Vol.I:135-136.

Papaioannou, D.S.; Kyriakis, S.C.; Papasteriadis, A.; Roumbies, N.; Yannakopoulos, A. and Alexopoulos, C. (2002) Effect of in-feed inclusion of natural zeolite (clinoptilolite) on certain vitamin, macro and trace element concentrations in the blood, liver and kidney tissues of sows. Res. Vet. Sci., 72(1):61-8.

Papantoniou,N.;Ismailos,G.;Daskalakis,G.;Karabinas,C.; Mesogitis ,S.; papapanagiotou,A.; Antsaklis,A. (2007) (Pharmacokinetics of oral cefatrizine in pregnant and non –pregnant women with reference to fetal distribution .) Fetal Diagn Ther.22(2):100-6.

Pastor L, del Olmo L, Lorenzo C, Almaraz A, Belmonte A, Coca MC, Caro-Patón A. (2002) [Effects of oxytetracycline on rat liver]. Gastroenterol Hepatol. 25(9):529-33.

Paul,J. ;Conkie, D. and Freshney, R.I.(1969) Erythropoietic cellpopulation changes during the hepatic phase of erythropoiesis in the foetal mouse. Cell Tissue Kinet. 2:283-294.

Penna, SC., Medeiros, MV., Aimbre, FS., Faria-Neto, HC., Sertie, JA and Lopes-Martins, RA. (2003) Anti- inflammatory effect of the hydral- coholic extract of *Zingiber officinal* rhizomes on rat paw and skin edema. Phytomed; 10: 381- 385 .

Philips, S. (1993) *Zingiber officinal* (ginger)-an antiemetic for day case surgery. Anaesthesia; 48: 715 - 7 .

Pieroni A, Torry B. (2007) Does the taste matter? Taste and medicinal perceptions associated with five selected herbal drugs among three ethnic groups in West Yorkshire, Northern England. J Ethnobiol Ethnomed.3:21.

Pindborg, JJ. (1982) Aetiology of developmental enamel defects not related to fluorosis. Int Dent J. 32(2):123-34.

Portnoi G, Chng LA, Karimi-Tabesh L, Koren G, Tan MP, Einarson A. (2003) Prospective comparative study of the safety and effectiveness of ginger for the treatment of nausea and vomiting in pregnancy. Am J Obstet Gynecol. 189(5):1374-1377.

Portnoi, G.; Chng, L.A.; Karimi-Tabesh, L.; Koren, G.; Tan, M.P. and Einarson, A. (2004) Prospective comparative study of the safety and effectiveness of ginger for the treatment of nausea and vomiting in pregnancy. Am. J. Obstet Gynecol. 189(5):1374-7.

Pride,G.;Cleary,R.and Hamburger,R. (1973) (Disseminated intravascular coagulation associated with tetracycline-induced hepatorenal failure during pregnancy)American Journal of obstetrics.585-586.

Q

Quian, DS, and Liu, ZS. (1992) Pharmacologic studies of antimotion sickness actions of ginger. Chung Kuo Chung His Chieh Ho Tsa Chih;12:95–8.

R

Raddy, AC and Lokesh, BR. (1992) Studies on spice principles as antioxidants in the inhibition of lipid peroxidation of rat liver microsomes. Mol. Cell. Biochem., 111 (1-2): 117 – 24.

Reboud AM, Dubost S, Reboud JP. (1982) Photoincorporation of tetracycline into rat-liver ribosomes and subunits. Eur J Biochem. 124(2):389-96.

Reynold, E.S. (1963) Theuse of lead citrate at high Phase an electronopaque stain in electron microscopy. J. Cell Biol., (17):208-212.

Rhode J, Fogoros S, Zick S, Wahl H, Griffith KA, Huang J, Liu JR. (2007) Ginger inhibits cell growth and modulates angiogenic factors in ovarian cancer cells. BMC Complement Altern Med. 7:44.

Richard,A.;Zager,Z. andJohnson,J. (2001) (Renal cortical cholesterol accumulation is an integral component of the systemic stress response). Kidney International ,Vol.60 pp.2299-2310.

Roberts AT, Martin CK, Liu Z, Amen RJ, Woltering EA, Rood JC, Caruso MK, Yu Y, Xie H, Greenway FL. (2007) The safety and efficacy of a dietary herbal supplement and gallic acid for weight loss. J Med Food.;10(1):184-8.

Romanowska, J.; Kifer-Wysocka, E. and Matysiak, W., (1991) [Histochemical picture of kidneys from pregnant female rats subjected to the action of vibramycin] .Ann Univ Mariae Curie Skłodowska[Med] , 46:149-54.

Romer, A.S. (1968) The vertebrate body. Third Edition. Published by W. B Saunders Company, Philadelphia, London. P:358-361.

Rømert P, Matthiessen ME. (1986) Tetracycline-induced changes in hepatocytes of mini-pigs and mini-pig foetuses as revealed by electron microscopy. Acta Pathol Microbiol Immunol Scand A. 94(2):125-31.

Romm, A., Herbalist, MD. Mary L., Hardy, MD. and Mills, S. (2010) Ginger. Botanical Medicine for Women's Health. P:536-538.

Rong, X., Peng, G., Suzuki, T., Yang, Q., Yamahara, J. and Li, Y. (2009) Regulatory Toxicology and Pharmacology. Volume 54, Issue 2, P:118-123.

Rothman KF, Pochi PE. (1988) Use of oral and topical agents for acne in pregnancy. J Am Acad Dermatol. 19(3):431-42.

Sabina EP, Rasool M, Mathew L, Ezilrani P, Indu H. (2010) 6-Shogaol inhibits monosodium urate crystal-induced inflammation-an in vivo and in vitro study. *Food Chem Toxicol.* 48(1):229-35.

Sadler, T.W. (2000) (Langnan, s. Medical Embryology) 8th Edition Lippincot Williamsauq Wilkins. New York, London, Tokyo.

Saito S, Ishikawa H, Ohtani M, Kawai K, Miyanaga N, Koiso K.(1990) [Basic study of nephrotoxicity of antibiotics. I. Studies of the effects of antibiotics on nucleic acids and protein metabolism in rat kidneys]. *Nippon Hinyokika Gakkai Zasshi.* 81(2):275-81.

Sakr, S.A., (1999). Pyrethroid inhalation induced hepatotoxicity in albino rats. *Oxford Res. Forum J.*, 1(1): 27-31.

Sakr,S. A. (2007) Ameliorative Effect of Ginger (*Zingiber officinale*) on Mancozeb Fungicide Induced Liver Injury in Albino Rats. *Australian Journal of Basic and Applied Sciences*, 1(4): 650-656.

Sambaiah K, Srinivasan K. (1991) Effect of cumin, cinnamon, ginger, mustard and tamarind in induced hypercholesterolemic rats. *Nahrung.* 35(1):47-51.

Sang S, Hong J, Wu H, Liu J, Yang CS, Pan MH, Badmaev V, Ho CT. (2009) Increased growth inhibitory effects on human cancer cells and anti-inflammatory potency of shogaols from *Zingiber officinale* relative to gingerols. *J Agric Food Chem.* 57(22):10645-50.

Sanjay, P.A.; Santosh, L.V. and Ramesh, K.G., (2004) Anti-diabetic activity of *Zingiber officinale* in streptoztocrin-induced type 1 diabetic rats. *J. Pharmacy and Pharmacology.*, 56:101-105.

Sattler,F.R.;Weitekam,M.R. and Ballard,J.O.(1986a) Potential for bleeding with the new Beta-lactam antibiotics. *Ann. Intern.Med.* 105:924-931.

Sattler,M.D.;Michael,R.;Weitekamp ,M.D. and James,o.(1986b) (Potential for Bleeding with the new Beta-lactam Antibiotics).*Annals of Internal Medicin.*926-931.

Schmucker, D. L. (1990) Hepatocyte fine structure during maturation and senescence. J. Electron. Microsc. Tech. 14(2):106-125.

Schultz jc, Adamson JS Jr, Workman WW, Norman TD. (1963) fatal liver disease after intravenous administration of tetracycline in high dosage. N Engl J Med. 269:999-1004.

Serreau R. (2010) [Drugs during preeclampsia. Fetal risks and pharmacology]. Ann Fr Anesth Reanim. 29(4):e37-46.

Shaker, R. and El-Baz, N. (2001) Experimenatal study on the effect of Tramadol Hydrochloricde (Central Analgesic) on the kidney of Adult Albino rat. Sc. J. Med., Vol.22.No. 1, pp:179-192.

Shanmugam KR, Ramakrishna CH, Mallikarjuna K, Reddy KS. (2010) Protective effect of ginger against alcohol-induced renal damage and antioxidant enzymes in male albino rats. Indian J Exp Biol. 48(2):143-9.

Shanmugasundaram KR, Ramanujam S, Shanmugasundaram ER. (2010) Amrita Bindu--a salt-spice-herbal health food supplement for the prevention of nitrosamine induced depletion of antioxidants. J Ethnopharmacol. 42(2):83-93.

Sharma, JN., Srivastava, KC and Gan, EK. (1994) Suppressive effects of eugenol and ginger oil on arthritic rats. Pharmacology; 49:314- 318.

Shauer BA, Lukacs L, Zimmerman HJ. (1974) Biochemical indices of tetracycline hepatic injury in rats. Proc Soc Exp Biol Med. 147(3):868-72.

Shcherbakova,E.G.; Rastunova,G.A.; Medved- eva,MM.; Zhuravieva,T.P.; Kuchuskev,S.H. A.; Lazareva,D.N. and Sobolev,v.R.(1977) Effect of gentamicin on the state Of cell resistance.J.Antibiotiki (Mosc.),22(5):413-417.

Shimoda H, Shan SJ, Tanaka J, Seki A, Seo JW, Kasajima N, Tamura S, Ke Y, Murakami N. (2010) Anti-inflammatory properties of red ginger (*Zingiber officinale* var. *Rubra*) extract and suppression of nitric oxide production by its constituents. J Med Food. 13(1):156-62.

Shosyrev, A.M. (1989) Comparative Stimulation of the embryotoxic action of some Antibiotics. *anion Res. Chem pharm .Inst. Moscow ,ussR.* 34 (10):779-782.

Siddaraju, M.N. and Dharmesh, S.M. (2007) Inhibition of gastric H+, k+, ATPase and Helicobacter pylori growth by phenolic antioxidants of *Zingiber officinale*. *Mol. Nutr. Food. Res.*, 51(3): 324-332.

Silva, J.R. and Rocha, R.F. (1995) Efeitos do cloridrato de tetraciclina em fetos de ratas tratadas com o antibiotico. *Rev. Odontol. UNESP*, 24: 109-115.

Singh Y, Garg DN, Kapoor PK, Mahajan SK. (2004) Isolation of Mycoplasma bovoculi from genitally diseased bovines and its experimental pathogenicity in pregnant guinea pigs. *Indian J Exp Biol.* 42(9):933-6.

Skosyrera,A.M.;Akhtamova ,Z.M.; Golovanova, G.V.and Gavorskii,A.(1986) (Effect of cefuroxime on embryonic and fetal development in experimental .*Antibiot.Med.Biotekhol*,31;277-280.

Skosyreva, A.M. (1989) [Comparative evaluation of the embryotoxic effect of various antibiotics]. *Antibiot Khimoter.*, 34(10): 779-82.

Slonitkaia, N.N. and Mikhalets, G.A. (1975) Toxic action of morphocycline on pregnancy rats. *Antibiotiki.*, 20(2):158-61.

Solera, J. (2000) Treatment of human brucellosis. *J Med Liban.* 48(4):255-63.

Srinivasan K, Sambaiah K. (1991) The effect of spices on cholesterol 7 alpha-hydroxylase activity and on serum and hepatic cholesterol levels in the rat. *Int J Vitam Nutr Res.* 61(4):364-9.

Srivastara, KC and Mostafa, T. (1992) Ginger (*Zingiber officinal*) in rheumatism and musculoskeletal disorders. *Med. Hypothesis*; 39 (4): 342 - 8 .

Standiford, H.C. (2000) Tetracyclines and chloramphenicol. In, Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases, 5th ed. (Mandell, G.L. Bennett, J.E., and Dolin, R., eds) Churchill Livingstone, Philadelphia, pp.336-348.

Stevens,A.;Lowe ,J.and young ,B.(2003) (Wheaters Basic histopathology). 4th Ed louis , Baltimor, Toronto .pp:33.

Suekawa,M. Ishige,A. Yuasa, K. Sudo,K. Aburada , M. and Hosoya,E. (1984) Pharmacological studies on ginger. I. Pharmacological actions of pungent constituents, (6)-gingerol and (6)-shogaol. J Pharmacobiodyn 7, pp. 836-848.

Summ HD, Draeger E, von Wasielewski E. (1976) [On the inhibitory effect of chloramphenicol on mitochondrial protein synthesis as a possible cause of its selective toxic side effects (author's transl)]. Arzneimittelforschung. 26(1):28-32.

Surh, Y. (1999). Molecular Mechanisms of Chemopreventive Effects of Selected Dietary and Medicinal Phenolic Substances, Mutat. Res 428(1-2): 305-327.

Sutton TA, Kelly KJ, Mang HE, Plotkin Z, Sandoval RM, Dagher PC. (2004) Minocycline reduces renal microvascular leakage in a rat model of ischemic renal injury. Am J Physiol Renal Physiol. 288(1):F91-7.

Suzuki ,H.;Yagi,M.;Saito,K.and Zuzuki,K.(2007) (Embryonic pathogenesis of hypogonadism and renal hypoplasia in hgn/hgn rats characterized by male sterility ,reduced female fertility and progressive renal insufficiency). Congenital Anomalies ;47,34-44.

Swanson-Biearman B, Dean BS, Lopez G, Krenzelok EP. (1988) The effects of penicillin and cephalosporin ingestions in children less than six years of age. Vet Hum Toxicol;30:66-67.

Sweet ,D.H .(2004) (Organic anion transporter(Slc22a) family members as Mediators of toxicity) Toxicology and Applied pharmacology 198-215.

T

Taber,S.S.;Mueller,B.A.(2006) (Drug-associated renal dysfunction). Crit care Clin. 22(2):357-74.

Tanabe, M., Chen, YD., Saito, K and Kano, Y. (1993) Cholesterol biosynthesis inhibitory component from *Zingiber officinal* Roscoe. Chem Pharm Bull (Tokyo); 41 (4) : 710 - 3.

Tang, W. and Eisenbrand, G. (1992) Chinese drugs of plant origin. Chemistry, pharmacology, and use in traditional and modern medicine, Springer-Verlag, Berlin.

Taylor,P.(1986) Handling : the reproductive cycle and mating In: Practical Teratology ,A cademic Press.Inc.London,Copyright C, PP:3-9.

Terao , K. and Miyaki, K. (1968) Different Susceptibilities of Chick Embryo Liver Cells in vitro to Aflatoxin, Actinomycin D, and Mitomycin C. Zeitschrift fiir Krebsforschung, 71:199-207.

Teslina OV, Anufrieva RG, Zel'tser IZ, Balabanova EL, Lapchinskaia AV. (1975) [Experimental study of the embryotropic action of tetracycline]. Antibiotiki. (9):839-42.

Thatte U, Bagadey S, Dahanukar S. (2000) Modulation of programmed cell death by medicinal plants. Cell Mol Biol (Noisy-le-grand). 46(1):199-214.

Thomson M, Al-Qattan KK, Al-Sawan SM, Alnaqeeb MA, Khan I, Ali M. (2002) The use of ginger (*Zingiber officinale* Rosc.) as a potential anti-inflammatory and antithrombotic agent. Prostaglandins Leukot Essent Fatty Acids. 67(6):475-8.

Tiran, D. (2003) The use of herbs by pregnant and childbearing women: a risk–benefit assessment. Complementary Therapies in Nursing & Midwifery . 9, 176–181.

Tolins,J.and Raij,L.(1990) (Chronic (Amphotericin B) Nephrotoxicity in the Rat protective effect of Calcium Channel Blockade). Journal of the American Society of Nephrology. 98-102.

Townson, S.; Tagboto, S.; McGarry, H.F.; Egerton, G.L and Taylor, M.J., (2006) Onchocerca parasites and *Wolbachia*

endosymbionts: evaluation of a spectrum of antibiotic types for activity against *Onchocerca gutturosa* *in vitro*. Filaria J. 5: 4.

Tsuda S, Murakami M, Matsusaka N, Kano K, Taniguchi K, Sasaki YF. (2001) DNA damage induced by red food dyes orally administered to pregnant and male mice. Toxicol Sci. 61(1):92-9.

Tsukahara K, Nakagawa H, Moriwaki S, Takema Y, Fujimura T, Imokawa G. (2006) Inhibition of ultraviolet-B-induced wrinkle formation by an elastase-inhibiting herbal extract: implication for the mechanism underlying elastase-associated wrinkles. Int J Dermatol. 45(4):460-8.

Trachtman H, Futterweit S, Greenwald R, Moak S, Singhal P, Franki N, Amin AR. (1996) Chemically modified tetracyclines inhibit inducible nitric oxide synthase expression and nitric oxide production in cultured rat mesangial cells. Biochem Biophys Res Commun. 229(1):243-8.

Triggs CM, Munday K, Hu R, Fraser AG, Gearry RB, Barclay ML, Ferguson LR. (2010) Dietary factors in chronic inflammation: Food tolerances and intolerances of a New Zealand Caucasian Crohn's disease population. Mutat Res.

Tripathi S, Bruch D, Kittur DS. (2008) Ginger extract inhibits LPS induced macrophage activation and function. BMC Complement Altern Med. 8:1.

Tripathi S, Maier KG, Bruch D, Kittur DS. (2007) Effect of 6-gingerol on pro-inflammatory cytokine production and costimulatory molecule expression in murine peritoneal macrophages. J Surg Res. 138(2):209-13.

Tune ,B.M.(1975) (Relationship between the transport and toxicity of Cephalosporins in the kidney). J.Infect Dis. 132:189-194.

Tune,B.M.(1994) (Renal tubular transport and nephrotoxicity of B-lactam antibiotics:structure -activity relationships .Minor Electrolyte Metab .20:221-231.

U

Uz E, Karatas OF, Mete E, Bayrak R, Bayrak O, Atmaca AF, Atis O, Yildirim ME, Akcay A. (2009) The effect of dietary ginger (*Zingiber officinale Rosc*) on renal ischemia/reperfusion injury in rat kidneys. *Ren Fail.* 31(4):251-60.

V

Van den Anker,J.N.;Hop ,W.C.;Schoemaker,R.C.and Neijens, H.J. (1995) (Cefazidime pharmacokinetics in preterm infants : effect of postnatal age an d postnatal exposure to indomethacin). *J Clin pharmacol.* 40(5) 439-43.

Van-den-Bogert, C. Blaauw, E. H., Dontje, B.H., Hulstaert, C.E., Hardonk, M.J. and Kroon, A.M. (1983) The effect of doxycycline on polyvinylpyrrolidone-induced granuloma. *Virchows, Arch. [B].* 51: 39-50.

Verma RJ, Asnani V. (2007) Ginger extract ameliorates paraben induced biochemical changes in liver and kidney of mice. *Acta Pol Pharm.* 2007 May-Jun;64(3):217-20.

Verma, S.K. Singh, J. Khamesra , R. and Bordia, A. (1993) Effect of ginger on platelet aggregation in man. *Indian J Med Res* 98, pp. 240–242.

Verma,S.and Kieff,E. (1975) (Cephalexin-related nephropathy .*J.A.M. A.*234:618-619.

Vijaya Padma V, Arul Diana Christie S, Ramkuma KM. (2007) Induction of apoptosis by ginger in HEp-2 cell line is mediated by reactive oxygen species. *Basic Clin Pharmacol Toxicol.* 100(5):302-7.

Vlastarakos PV, Nikolopoulos TP, Manolopoulos L, Ferekidis E, Kreatsas G. (2008) Treating common ear problems in pregnancy: what is safe? *Eur Arch Otorhinolaryngol.* 265(2):139-45.

Vutyavanich, T., Kraisin, T. and Ruangsri, R. (2001) Ginger for nausea and vomiting in pregnancy: Randomized, double-masked, placebo-controlled trial. *Obstetrics & Gynecology*, Volume 97, Issue 4, P:577-582.

W

Wang J, Wei Q, Wang CY, Hill WD, Hess DC, Dong Z. (2004) Minocycline up-regulates Bcl-2 and protects against cell death in mitochondria. J Biol Chem. 279(19):19948-54.

Watanabe, H.; Sakai, T.; Kobayashi, N.; Fukuday and Yabutak (1996) Glomerular basement membrane outpockets and glomular growth in the postnatal development of the rat kidney. Pediatr. Nephrol., 10(4): 461-6.

Watt G, Kantipong P, Jongsakul K, Watcharapichat P, Phulsuksombati D. (1999) Azithromycin activities against Orientia tsutsugamushi strains isolated in cases of scrub typhus in Northern Thailand. Antimicrob Agents Chemother. 43(11):2817-8.

Weidner MS, Sigwart K. (2000) The safety of ginger extract in the rat. J Ethnopharmacol. 73(3):513-520.

Weidner, M.S. and Sigwart, K.(2001) Investigation of the teratogenic potential of a zingiber officinale extract in the rat. Reprod Toxicol .,15(1):75-80.

Weng CJ, Wu CF, Huang HW, Ho CT, Yen GC. (2010) Anti-invasion effects of 6-shogaol and 6-gingerol, two active components in ginger, on human hepatocarcinoma cells. Nano Lett. [Epub ahead of print].

Westphal JF, Vetter D, Brogard JM. (1994) Hepatic side-effects of antibiotics. J Antimicrob Chemother. 33(3):387-401.

Wilkinson, J. M., (2000a) Effect of ginger tea on the fetal development of Sprague-Dawley rats. Reprod Toxicol., 14(6):507-12.

Wilkinson, J. M.(2000b) What do we know about herbal morning sickness treatments? - a review of the literature. Midwifery; 16: In press.

Wilson WR. (1977) Tetracyclines, chloramphenicol, erythromycin, and clindamycin. Mayo Clin Proc. 52(10):635-40.

Wilson WR, Cockerill FR 3rd. (1983) Tetracyclines, chloramphenicol, erythromycin, and clindamycin. Mayo Clin Proc. 58(2):92-8.

Wiśniewska IE, Gawlik Z. (1982) The effect of doxycycline upon alkaline phosphatase activity in rat kidney (quantitative study using the interferometric technique). Folia Histochem Cytochem (Krakow). 20(3-4):157-62.

Woolhouse, M. (2006) Complementary medicine for pregnancy complications. Aust Fam Physician. 35(9):695.

Worret WI, Fluhr JW. (2006) [Acne therapy with topical benzoyl peroxide, antibiotics and azelaic acid]. J Dtsch Dermatol Ges. 4(4):293-300.

Wruble,G. D. Ladman,A.J. Britt, G.G. and Cummins,A. J. (1965), J. Am. Med. Ass. 192, 6.

Y

Yagihashi S, Miura Y, Yagasaki K. (2008) Inhibitory effect of gingerol on the proliferation and invasion of hepatoma cells in culture. Cytotechnology. 57(2):129-36.

Yamahara, J., Mochizuki, M., Rong, HK., Matsuda, H and Fujimura, H. (1988) The anti-Ulcer effect in rats of ginger constituents. Journal of Ethnopharmacology; 23: 299 – 304.

Yaman I, and Balikci E. (2010) Protective effects of nigella sativa against gentamicin-induced nephrotoxicity in rats. Exp Toxicol Pathol. 62(2):183-90.

Yang G, Zhong L, Jiang L, Geng C, Cao J, Sun X, Ma Y. (2010) Genotoxic effect of 6-gingerol on human hepatoma G2 cells. Chem Biol Interact. 185(1):12-7.

Yao H, Tong J, Wang Z, Zhang ZQ, Chen JX. (2007) [Effect of acupoint application therapy on expression of IL-4 and IFN-gamma in lung tissue of asthma rats]. Zhen Ci Yan Jiu. 32(3):174-8.

Yasui T, Fujita K, Asai K, Kohri K. (2002) Osteopontin regulates adhesion of calcium oxalate crystals to renal epithelial cells. Int J Urol. 9(2):100-8.

Yemitan, O.K. and Izegbu, M.C. (2006) Protective effects of Zingiber officinale (Zingiberaceae) against carbontetrachloride and acetaminophen-induced hepatotoxicity in rats. Phytother Res., 20(11):997-1002.

Yong Jung,K.J.;Michio,T.;Minoru,S.;Shinichi,N.and Akihiro, T.(2002) (Involvement of rat organic anion transport 3 (rOAT3) in cephaloridine -induced nephrotoxicity : In comparison with rOAT1) Life Sciences 70 .1861-1874.

York DA, Thomas S, Greenway FL, Liu Z, Rood JC. (2007) Effect of an herbal extract Number Ten (NT) on body weight in rats. Chin Med. 2:10.

Yusof YA, Ahmad N, Das S, Sulaiman S, Murad NA. (2008) Chemopreventive efficacy of ginger (Zingiber officinale) in ethionine induced rat hepatocarcinogenesis. Afr J Tradit Complement Altern Med. 6(1):87-93.

Z

Zakeri, B. and Wright, G. D., (2008) Chemical biology of tetracycline antibiotics. Biochem. Cell Biol. 86, 124-136.

Zítková L, Stastná J, Dobrovský K, Tousek J. (1982) Causes for rifampicin toxicity--experimental study. Czech Med. 5(4):210-7.

Zussman, w. v. (1968) Hepatic alterations following experimental tetracycline toxicity in rats. Anat. Rec. 162, 301-312.